arXiv:1611.02527v1 [math.LO] 8 Nov 2016

WEAK KÖNIG'S LEMMA IMPLIES THE UNIFORM CONTINUITY THEOREM: A DIRECT PROOF

MATTHEW HENDTLASS

Abstract. We show in Bishop's constructive mathematics—in particular, using countable choice—that weak König's lemma implies the uniform continuity theorem.

In [1] Hannes Diener proved, as part of the programme of reverse constructive mathematics, that weak König's lemma

WKL: Every infinite, decidable, binary tree has an infinite path.

implies the uniform continuity theorem

UCT: Every pointwise continuous function $f : [0, 1] \to \mathbf{R}$ is uniformly continuous.

in Bishop's constructive mathematics. Diener's proof relies on several other results in reverse constructive mathematics; we give a short, direct proof.

Theorem 1. *Weak König's lemma implies the uniform continuity theorem.*

The idea of our proof is simple: given a continuous function $f : [0, 1] \to \mathbf{R}$ and some $\varepsilon > 0$, we use WKL to focus in on a point where the function exhibits (almost) greatest variation within ε and then use continuity at that point to find our modulus δ of uniform continuity for ε. Before proving Theorem 1 we must set up some notation.

For each $n > 0$ we let $S_n = \{0, 2^{-n}, \ldots, 1 - 2^{-n}, 1\}$ and we write $\mathcal{D}$ for the set $\cup\{S_n : n \in \omega\}$ of dyadic rationals. We define a one-one function g from the set $2^{<\omega}$ of finite binary sequences to $\mathcal{D}$ by

$$g(a) = \sum_{i=0}^{|a|-1} a(i)2^{-(i+1)}$$

and associate $x \in S_n$ with the unique finite binary string a of length n such that $g(a) = x$. The *sum* of two finite binary sequences a, b each of length n is the binary sequence $a \oplus b$ of length $2n$ given by

$$a \oplus b(n) = \begin{cases} a(n/2) & n \text{ is even} \\ b((n-1)/2) & n \text{ is odd.} \end{cases}$$

Both g and $\oplus$ extend to functions on infinite binary sequences and we make no notational distinction between the functions on finite and infinite sequences. We let π_0, π_1 be the left and right inverses of $\oplus$ respectively; that is $\pi_0(a \oplus b) = a$ and $\pi_1(a \oplus b) = b$ for all suitable pairs $a, b \in 2^{<\omega} \cup 2^{\omega}$. For $\alpha \in 2^{\omega}$ and $n \in \mathbf{N}$, $\bar{\alpha}(n)$ denotes the unique binary sequence of length n that α extends. The downward closure of subset A of $2^{<\omega}$ is $A\downarrow = \{a \in 2^{<\omega} : \exists_{a' \in A}(a < a')\}$, where $a < a'$ if a' extends a.

For a given function $f : [0, 1] \to \mathbf{R}$ we define a predicate φ_f on $\mathbf{R}^+ \times \mathbf{R}^+$ by

$$\varphi_f(\varepsilon, \delta) \equiv \exists_{x,y \in [a,b]}(|x - y| < \delta \wedge |f(x) - f(y)| > \varepsilon);$$

$\varphi_f(\varepsilon, \delta)$ holds if we have a witness that δ is not a modulus of uniform continuity for f, ε. Thus to show that $f : [0, 1] \to \mathbf{R}$ is uniformly continuous we must, given any $\varepsilon > 0$, find some $\delta > 0$ such

that $\varphi_f(\varepsilon,\delta)$ is false. The next lemma shows how we can use **WKL** to reduce the truth of $\varphi_f(\varepsilon,\delta)$ to whether or not some specific $x,y\in[0,1]$ are witnesses of $\varphi_f(\varepsilon,\delta)$.

Lemma 2. ***WKL*** $\vdash$ *Let $f:[0,1]\to\mathbf{R}$ be pointwise continuous and let δ,ε be positive real numbers. Then there exist $x,y\in[a,b]$ such that if δ is not a modulus of uniform continuity for f, then x,y witness this:*

$$\varphi_f(\varepsilon,\delta)\to|x-y|<\delta\wedge|f(x)-f(y)|>\varepsilon.$$

Proof. Using countable choice, construct a function $\gamma:\mathcal{D}^2\times\mathbf{N}\to 2$ such that

$$\begin{aligned}\gamma(x,y,n)=0 &\Rightarrow |x-y|>\delta-2^{-n}\vee|f(x)-f(y)|<\varepsilon+2^{-n},\\ \gamma(x,y,n)=1 &\Rightarrow |x-y|<\delta\wedge|f(x)-f(y)|>\varepsilon;\end{aligned}$$

further we may assume that γ is non-decreasing in the third argument. So if $\gamma(x,y,n)=1$, then x,y are witnesses of $\varphi_f(\varepsilon,\delta)$. Using γ, we can construct an increasing binary sequence $(\lambda_n)_{n\in\mathbf{N}}$ such that

$$\begin{aligned}\lambda_n=0 &\Rightarrow \forall_{x,y\in S_n}\gamma(x,y,n)=0,\\ \lambda_n=1 &\Rightarrow \exists_{x,y\in S_n}\gamma(x,y,m)=1.\end{aligned}$$

Finally we construct a decidable binary tree T as follows. If $\lambda_n=0$ we let $T_n=2^n$, and if $\lambda_{n-1}=1$ we set $T_n=\{\sigma*0:\sigma\in T_{n-1}\text{ and }|\sigma|=\mathrm{ht}(T)\}$. If $\lambda_n=1-\lambda_{n-1}$, we let x,y be the minimal elements of S_n such that $\gamma(x,y,n)=1$ and we set $T_n=(2^n\cup\{x\oplus y\})\downarrow$—the branch $x\oplus y$ is the unique branch of T_n with length $\mathrm{ht}(T_n)$, and it codes the witnesses x,y that δ is not a modulus of uniform continuity for ε. Then

$$T=\bigcup_{n\in\mathbf{N}}T_n$$

is an infinite decidable tree.

Using **WKL** we can construct an infinite path α through T. Set $x=g(\pi_0\alpha),y=g(\pi_1\alpha)$. Suppose there exist $u,v\in[0,1]$ such that $|u-v|<\delta$ and $|f(u)-f(v)|>\varepsilon$. Since f is pointwise continuous, there must exist such $u,v\in\mathcal{D}$. Hence $\gamma(u,v,n)=1$ for some $n\in\mathbf{N}$ such that $u,v\in S_n$, so $\lambda_n=1$. It now follows from the construction of T that x,y have the desired property. □

We recall a result of Hajime Ishihara [2]: **WKL** is equivalent to the longest path principle

> **LPP**: Let T be a decidable tree. Then there exists $\alpha\in 2^\omega$ such that for all n, if $\bar{\alpha}(n)\notin T$, then $T\subset 2^{<n}$.

To get a longest path for a decidable tree T apply **WKL** to the decidable tree

$$\{a\in 2^{<\omega}:a\in T\text{ or }\exists_{b\in T}(|b|=\mathrm{ht}(T)\wedge a>b)\},$$

where $|b|$ is the length of b and $\mathrm{ht}(T)$ is the height of T.

Here then is our **proof of Theorem 1**:

Proof. Let $f:[0,1]\to\mathbf{R}$ be a pointwise continuous function and fix $\varepsilon>0$. We define a function J taking finite binary sequences to subintervals of $[0,1]$ inductively: $J_{()}=[0,1]$ and if $J_u=[p,q]$, then $J_{u*0}=[p,(p+q)/2]$ and $J_{u*1}=[(p+q)/2,q]$. By repeated application of the lemma, let $x_u,y_u\in J_u$ be such that

$$\varphi_{f|_{J_u}}(\varepsilon,2^{-|u|})\to|x_u-y_u|<2^{-|u|}\wedge|f(x_u)-f(y_u)|>\varepsilon/2,$$

and using countable choice construct a decidable tree T such that

$$\begin{aligned}u\in T &\Rightarrow |f(x_u)-f(y_u)|>\varepsilon/2-2^{-|u|}\\ u\notin T &\Rightarrow |f(x_u)-f(y_u)|<\varepsilon/2.\end{aligned}$$

Let α be a longest path of T, and let ξ be the unique element of

$$\bigcap_{n\in\mathbf{N}} J_{\overline{\alpha}(n)}.$$

Using the continuity of f at ξ we can find $\delta > 0$ such that $|f(x) - f(y)| < \varepsilon/2$ for all $x, y \in (\xi - \delta, \xi + \delta)$; let n be such that $2^{-n+1} < \max\{\delta, \varepsilon\}$. If $u = \overline{\alpha}(n) \in T$, then $|x_u - y_u| < \delta$ and $|f(x_u) - f(y_u)| > \varepsilon - 2^{-n} > \varepsilon/2$ contradicting our choice of δ. Hence $\overline{\alpha}(n) \notin T$, so $T \subset 2^{<n}$. It follows from Lemma 2 and the construction of T that for all $x, y \in [0, 1]$, if $|x - y| < 2^{-n}$, then $|f(x) - f(y)| < \varepsilon$. □

References

[1] H. Diener, Weak König's lemma implies the uniform continuity theorem, Computability 2(1), p. 9–13, 2013.
[2] H. Ishihara, 'Weak König's Lemma Implies Brouwer's Fan Theorem: A Direct Proof', Notre Dame Journal of Formal Logic 47(2), p. 249–252, 2006.

School of Mathematics and Statistics, University of Canterbury, Christchurch 8041, New Zealand
E-mail address: matthew.hendtlass@canterbury.ac.nz